

Neslihan CUHACI<sup>1</sup>, Berna OGMEN<sup>2</sup>, Ali TAM<sup>1</sup>, Cevdet AYDIN<sup>1</sup>, Oya TOPALOGLU<sup>1</sup>, Aylin KILIC YAZGAN<sup>3</sup>, Gurkan DUMLU<sup>4</sup>, Reyhan ERSOY<sup>1</sup>, Bekir CAKIR<sup>1</sup>

<sup>1</sup>Yildirim Beyazıt University, Faculty of Medicine, Ataturk Education and Research Hospital, Department of Endocrinology and Metabolism, Ankara, TURKEY

<sup>2</sup>Ataturk Education and Research Hospital, Department of Endocrinology and Metabolism, Ankara, TURKEY

<sup>3</sup>Ataturk Education and Research Hospital, Department of Pathology, Ankara, TURKEY

<sup>4</sup>Yildirim Beyazıt University, Faculty of Medicine, Ataturk Education and Research Hospital, Department of General Surgery, Ankara, TURKEY

## INTRODUCTION

➤ The increased rate of thyroid malignancy as well as incidental and subcentimeter thyroid nodules have been attributed to increasing use of high-resolution US which can detect the non-palpabl or subcentimeter (maximum diameter  $\leq 1$  cm) thyroid nodules.

➤ We aimed to evaluate the sonographic features of the thyroid nodules between  $\leq 1$  cm and  $> 1$ cm according the histopathology results and to determine the ultrasonographical predictive factors for malignancy and an ultrasonographic score according the sonographic features to avoid unnecessary fine-needle aspiration biopsy (FNAB).

## METHODS

➤ We retrospectively evaluated 2233 nodules of 1118 patients who underwent thyroidectomy.

➤ Predictive factors for distinguishing between malignant and benign histopathologic results according the ultrasonographic features were evaluated by multivariate logistic regression analysis.

➤ Multiple binary logistic regression with the forward logistic regression method was used to develop the formula for recommending sonographically guided biopsy.

➤ Among the 2233 nodules 337 nodules were in the  $\leq 1$  cm (group 1), 1896 were in the  $>1$ cm (group 2).

➤ According the histopathological results, in group 1; 173 nodules were in the benign, 164 nodule were in the malignant group. Whereas in group 2; 1423 nodules were in the benign, 473 nodules were in the malignant group.

➤ In group 1, AP/T  $\geq 1$ , the presence of microcalcification, macrocalcification and hypoechoic pattern were significantly higher in the malignant group and in group 2, the presence of microcalcification, macrocalcification, hypoechoic and iso-hypoechoic pattern, solid texture, peripheral and intranodular vascularization pattern were significantly higher in the malignant group.

➤ In group 1, the best ultrasonographic index score was found  $>2$ , whereas in group 2 the it was found  $>4$ .

**Table .** Index scores related with US features that can predict malignancy in nodules  $\leq 1$  cm and  $>1$  cm

	Nodule size	
	$\leq 1$ cm	$>1$ cm
<b>Ultrasonographic index score</b>		
Benign	2.09 $\pm$ 1.19	3.97 $\pm$ 1.46
Malignant	3.04 $\pm$ 1.06	4.95 $\pm$ 1.70
<b>ROC analysis</b>		
Area under the curve	0.722	0.665
95% Confidence interval	0.667-0.777	0.636-0.693
p-value	$<0.001$	$<0.001$
<b>The best cut-off point</b>	$>2$	$>4$
Sensitivity	68.6%	58.1%
Specifity	66.5%	66.6%
PPV	64.4%	36.7%
NPV	70.6%	82.7%

## CONCLUSION

➤ Our US scoring may lead to clinicians and surgeons to diagnose thyroid malignancy more accurately and to select the nodules for FNAB especially in subcentimeter nodules.