TRAIL downregulates Endothelin 1 and upregulates Superoxide Dismutase 1 in human aortic endothelial cells under proatherogenic conditions

Forde H^{1,2}, Davenport C^{1,2}, Cummins PM², Smith D¹

- 1. Beaumont & RCSI Medical school, Beaumont, Dublin 9
- 2. School of Endothelial biology, DCU, Glasnevin Dublin 9



OBJECTIVES

- Tumour necrosis factor related apoptosis inducing ligand (TRAIL) may be involved in the pathogenesis of cardiovascular disease
- In vivo findings suggest that TRAIL may exhibit vasoprotective effects on the endothelium, although the mechanism of TRAIL-mediated vasoprotection remains poorly understood^{1,2}
- Atherosclerosis is a chronic inflammatory condition and *in vitro* studies have demonstrated that endothelial cells exposed to disturbed/oscillatory flow adopt a proinflammatory phenotype³




Figure 1. ECs exposed to laminar flow





Figure 2. ECs exposed to oscillatory flow

The aim of this study was to characterise the protective effects of TRAIL on inflammatory gene expression from vascular endothelial cells under proatherogenic conditions

METHODS

- Human aortic endothelial cells (HAECs) were grown to confluency in standardised Promocell media
- HAECs were seeded onto 0.6mm µ-slide plate at a concentration of 200,000 cells per slide
- The slide was attached to the IBIDI_® µ-slide pump system (Image 1) which was connected to an air pressure control pump
- Cells were exposed to the conditions outlined below






Image 1. Fluidic unit from the IBIDI μ-slide pump system. A proatherogenic environment is simulated by applying oscillatory flow to the μ-slide by separating the switching times of the valves on the fluidic unit

Image 2. PCR array analysis on Real Time PCR system

RESULTS




Figure 3. Effect of TRAIL on HAEC gene expression under oscillatory flow. Histogram/table shows upregulatory (blue) and downregulatory (red) effects of TRAIL on HAEC genes under oscillatory flow. Housekeeping genes for normalization include B2 macroglobulin and arachidonate 5-lipoxygenase

DISCUSSION

- TRAIL upregulated a number of genes including SOD1, which encodes the enzyme Superoxide dismutase, involved in the breakdown of reactive oxygen species (ROS)
- TRAIL downregulated EDN1 and EDNRA, which encodes Endothelin 1 and the Endothelin receptor. These potent proinflammatory proteins can mediate ROS generation through increased NADPH oxidase activity⁴
- TRAIL also downregulated the adhesion molecule genes CDH5 and PECAM, which are involved in the promotion of atherogenesis
- In conclusion, TRAIL appears to exert protective anti-inflammatory effects, in a proatherogenic environment, partly through reduction of oxidative stress

REFERENCES

- Di Bartolo BA., Chan J., Bennett MR., Cartland S., Bao S., Tuch BE., Kavurma MM. (2011).
 TNF-related apoptosis-inducing ligand (TRAIL) protects against diabetes and atherosclerosis in Apoe-/- mice. *Diabetologia*, 54(12), 3157-3167
- . Watt V., Chamberlain J., Steiner T., Francis S., Crossman D. (2011). TRAIL attenuates the development of atherosclerosis in apolipoprotein E deficient mice. *Atherosclerosis*, 215(2), 348-354.
- 3. Malek AM., Alper SL., Izumo, S. (1999). Hemodynamic shear stress and its role in atherosclerosis. *Jama*, 282(21), 2035-2042.
- Dong F., Zhang X., Wold LÈ., Ren Q., Zhang Z., Ren, J. (2005). Endothelin-1 enhances oxidative stress, cell proliferation and reduces apoptosis in human umbilical vein endothelial cells: role of ETB receptor, NADPH oxidase and caveolin-1. British journal of pharmacology, 145(3), 323-333

